STUDIES REGARDING THE EFFECT OF STORING CEREALS AT LOW TEMPERATURES UPON THE SENSORY QUALITIES OF FOOD

S. HORINCAR¹, Eugenia PRICOP²

^{1,} Sanitary Veterinary and Food Safety Department,Galaţi,

 $e\hbox{-}mail: sor in. hor in car@yahoo.com$

^{2,} Biochemistry Department, Faculty of Food Science and Engineering, "Dunarea de Jos" University, Galați

This study is focused on the effect of the storage conditions on the sensory qualities of the cooked rice. From the data analyzed in the paper we could see that after four weeks the storage temperature has a significant effect on a series of sensory qualities as: sulphur smell, hardness (resistance to chewing) and cohesion of the grain mass. The sulphur smell decreases significantly when the storage temperature increases from 5 to 35°C. This can be explained through the volatilisation of the components that give this smell at the same time with the increase in temperature. The sulphur smell decreases significantly after twenty weeks of storage at all storage temperatures. For the samples examined after four weeks of storage, their resistance to chewing increases at the same time with the increase in storage temperature. The samples with a low chewing resistance also show an increase of grain cohesion. After twenty weeks of storage, the increase in the storage temperature causes a decrease of adhesion, adhesion at chewing and grain cohesion. Also we can notice an increase of a rancid smell.

Key words: cereals, conservation, low temperatures

Cultivated rice at present sustains two-thirds of the world's population. A small amount of the rice crop is used as ingredients in processed foods and as feed, but the bulk is consumed as cooked rice. This pattern of usage results in the need to store rice over varying periods. During storage, a number of physico-chemical and physiological changes occur, this is usually termed ageing [3]. Although the mechanism of rice ageing is not fully understood, the changes of pasting properties, colour, flavour, and composition of rice can be observed [7], which affects milling, cooking and eating quality [9,10]. The sensory qualities of cereal destined for human consumption have an influence on their economical value on the world market.

These sensory qualities are influenced by: the type of cereal, climatic conditions, conditions during harvesting and the operations after harvesting (drying, storage and grinding conditions)[2]. It is therefore important to understand the connection between the storage conditions and the final product, so that we can

coordinate our processes in the desired way. The specific objective of this study was to evaluate the effects of rough rice, storage temperature, and storage duration on the cooking properties of rice. This study is particularly focused on the effect of the storage conditions on the sensory qualities of the cooked rice.

MATERIAL AND METHOD

The rice used in the experiment is long grain type from the Cypress variety, with 20,5% humidity at harvest. Immediately after harvest, the rough rice samples were cleaned in a dockage tester and immediately air-dried at room temperature (»20°C) The moisture content of the rice was measured by drying duplicate samples for 24 hr in an air oven at 130°C [1]. The rice was prepared for storage and brought to a 12,5% humidity. The temperature levels used for storage were 5, 20 and 35°C, and the samples were analyzed after a period of 4 and 20 weeks. Milled rice was cooked for the minimum cooking time [4]. A team of nine tasters was present for the experiment. The intensity of each analyzed attribute was appreciated on a scale from 1 to 15. The reference intensity for each attribute was assured during the whole time of the experiment.

RESULTS AND DISCUSSIONS Table for the second state of the second size of the second size

Table 1
Effect of storage temperature on sensory quality of cooked rice after four weeks of storage

Sensory characteristics	Storage temperature		
	5 °C	20 °C	35 °C
General impression	4,37	4,25	4,19
Flavor of starch	3,57	3,53	3,49
Rancid smell	2,48	2,27	2,57
Sulfur odor	1,79	1,56	1,18
Strength	2,71	2,79	2,57
Roughness	2,82	2,95	2,96
Hardness	2,96	3,08	3,35
Adhesion to chew	2,88	2,78	2,85
Hygroscopicity	5,72	5,89	5,66
Cohesion	4,93	4,87	4,63

From the data analyzed in the paper we could see that after four weeks the storage temperature has a significant effect on a series of sensory qualities as: sulphur smell, hardness (resistance to chewing) and cohesion of the grain mass. The sulphur smell decreases significantly when the storage temperature increases from 5 to 35°C. This can be explained through the volatilisation of the components that give this smell at the same time with the increase in temperature.

Table 2
Effect of storage temperature on sensory quality of cooked rice after twenty weeks of storage

ory characteristics	Storage temperature		
	5 °C	20 °C	35 °C
General impression	4,22	4,36	4,57
Flavor of starch	3,65	3,57	3,77
Rancid smell	2,59	2,33	2,68
Sulfur odor	0,72	0,66	0,97
Strength	3,18	2,42	2,02
Roughness	2,91	2,75	2,98
Hardness	2,73	2,62	2,98
Adhesion to chew	3,55	3,04	2,69
Hygroscopicity	5,53	5,21	5,34
Cohesion	5,34	4,72	4,31

The sulphur smell decreases significantly after twenty weeks of storage at all storage temperatures. For the samples examined after four weeks of storage, their resistance to chewing increases at the same time with the increase in storage temperature. The samples with a low chewing resistance also show an increase of grain cohesion. After twenty weeks of storage, the increase in the storage temperature causes a decrease of adhesion, adhesion at chewing and grain cohesion. Also we can notice an increase of a rancid smell.

CONCLUSIONS

Cooked rice from freshly harvested rice is generally stickier than that from aged rice and is not preferred by people. The changes in rice properties during aging have been reported to depend upon variety and storage conditions[5]. The changes have been attributed to the changes in cell wall and protein, interaction between proteins and breakdown products of lipids oxidation, and starch-protein interaction [6]. Villareal et all reported a gradual decrease of acetic acid-soluble proteins during storage [11]. Champagne reported that the number of disulphide bonds and the average molecular weight of oryzenin (major protein in rice) increased during storage of rice grains [5] .Hamaker found that low molecular weight subunits of oryzenin decreased and high molecular weight subunits increased during storage.

The data presented in this paper confirm the fact that the storage temperature and time have a significant influence on some of the sensory properties of cereals. Therefore knowing the effects of the storage temperature and time can lead to an

improvement of these properties and it can avoid the degradation of the sensory properties of the stored products.

BIBLIOGRAPHY

- 1. Barna, O., 2007 Research on low temperature cereal preservation, PhD thesis University "Dunarea de Jos" of Galati, Romania.
- Barna, O., Tofan, I., 2001 Influence of temperature and duration of storage on technological properties of starch from rice. Newsletter for milling and baking industries vol. 13 (2):130-135.
- 3. Batcher, O.M., Helmintoller, K..F., 1956 Development and application of methods for evaluating cooking and eating quality of rice. Rice J. 59, 4-8, 32.
- Champagne, E.T., Lyon, B.G., Min, B.K., Vinyard, B.T., Bett, K.L., Barton, F.E., 1998 Effects of post harvest processing on texture profile analysis of cooked rice. Cereal
 Chem. 75: 181-186.
- 5. Chrastil, J., 1990 Chemical and physicochemical changes of rice during storage at different temperatures. J. Cereal Sci. 11:71-85.
- Chrastil, J. 1994 Effect of storage on the physicochemical properties and quality factors of rice. Pages 49-81 in: Rice Science and Technology. W. E. Marshall and J.I. Wadsworth, eds. Marcel-Dekker: New York.
- 7. Hamaker, B. R., Siebenmorgen, T. J., and Dilday, R. H., 1993 *Aging of rice in the first six months after harvest.* Ark. Farm Res. 42(1):8-9.
- 8. Indudhara, Swamy, Y.M., Sowbhagya, C.M., Bhattacharya, K.R., 1978 Changes in the physicochemical properties of rice with aging. J. Sci. Food Agric. 29:627-639.
- 9. Jindal, V.K., Siebenmorgen, T.J., 1987 Effects of oven drying temperature and drying time on rough rice moisture content determination. Trans. ASAE 30:1185-1192.
- 10. Villareal, R.M., Resurreccion, A.P., Suzuki, L.B., Juliano, B.O., 1976 Changes in the physicochemical properties of rice during storage. Starch/Staerke 28:88-94.
- 11. ***, 2000 AACC International. Approved Methods of the American Association of Cereal Chemists,10th Ed. Methods. The Association: St. Paul, MN.