

## RESISTANCE AND PRODUCTIVE QUALITIES OF PIGS WHEN USED RBS AND IMMUNOLAC

N. Cherniy<sup>1\*</sup>, Olga Machula<sup>1</sup>, Elena Tkachuk<sup>1</sup>, Valentina Rud<sup>2</sup>,  
Y. Schepetilnikov<sup>1</sup>

<sup>1</sup>Kharkov State Zooveterinary Academy, Kharkov, Ukraine

<sup>2</sup>Odessa State Agrarian University, Odessa, Ukraine

### Abstract

*The purpose of the research was to study the effect of stimulating drugs - the regenerating biological stimulant (RBS) and Immunolac on the resistance of pigs and their productive qualities. The hygienic, clinical, biochemical, immunological, veterinary, zootechnical methods and methods of variation statistics were used to reach the goal. The piglets of 1-60-day age of Large White breed and Landrace hybrids were used as an object for the experiment. The blood and blood serum, immunological (CIC, T- and -B-lymphocytes) humoral (BASK, LASK), cellular indices (FAN and FCH) and the parameters of live weight were the subject of the study. The choice of the object for the research was determined by the lack of the data on the use of RBS and Immunolac for the pigs and the effect of the above preparations on the health and immunological status of the pigs. By the results of the experiment it has been found out that when piglets are grown in the conditions of uncontrolled microclimate without the use of the drugs their genetic productive potential is not fully realized, the above preparations do not cause immunosuppression, they are harmless at doses of 0,05 ml/kg of body weight, they improve general condition of the pigs and do not cause any changes that indicate about pathological condition and growth depression.*

**Key words:** piglets, immunostimulants, globulins, resistance, live weight

### INTRODUCTION

Pork production can be effective only if hygienic requirements are met and the animals are provided with full-value feeds [8]. The appearance of stresses and the development of immunodeficiency in young animals as a result of the discrepancy between the physiological capabilities of the pigs and the biotic factors of the environment (indoor maintenance, fluctuations of temperature, high humidity and bacterial pollution of the air) that reduce the immune status of the piglets, especially in the early postnatal period interfere the introduction of the intensive technologies in many pig-breeding enterprises [4]. In this connection, one of the problems in pig - breeding [2] is firstly, the development of measures aimed at the reduction of the effect of the unfavorable microclimate factors on swine homeostasis,

secondly, the prevention of immunodeficiencies [1, 3] by the use of pro-and-prebiotics, immunostimulants [5,7] that lead to the considerable damage. In our experiment, we used the immunostimulants that increase the level of realization of the bioresource potential of pigs according to the instructions that led to the research. The purpose of the study was to study the effect of the immunostimulating drugs - RBS and immunolac on the immunological state and productive parameters of piglets. The regenerating biostimulator (RBS) is a complex of organic compounds of the animal tissue that stimulates nonspecific immunity, increases the protective functions of the body in relation to microbes and viruses, activates T- and B- lymphocytes. Immunolac is a product of enzymatic hydrolysis from the cell wall of *Lactobacillus*, it enhances hematopoiesis, activates cellular and humoral factors of nonspecific resistance.

---

\*Corresponding author: dmchorn@ukr.net

The manuscript was received: 19.09.2017

Accepted for publication: 14.02.2018

## MATERIALS AND METHODS

The research was carried out at the pig-breeding plant "Stepnoy", Zaporizhzhya region. According to the principle of analogues, three groups of 15 piglets of 1-day age piglets of Large White breed × Landrace were formed. The piglets of the control group were administered 0.9% sodium chloride solution at the dose of 1 ml / head, the piglets of the experimental group 1 received Immunolac at the dose of 0.05 ml/kg body weight, the piglets of the experimental group 2 received RBS at the dose of 0.05 ml/kg live weight on the 3<sup>rd</sup>, 7<sup>th</sup>, 15<sup>th</sup> day of life. The above drugs were administered intramuscularly at the indicated doses on days 3, 7, 15. During the experiment the sanitary and hygienic state in the sections was monitored for the parameters: temperature, relative humidity, air velocity, content of carbon dioxide, ammonia, the contamination of the air with microflora. The clinical and physiological state of the animals was assessed by the morphological parameters of the blood by counting erythrocytes and leukocytes in the Goryaev's chamber with net. The phagocytic activity of neutrophils was determined in relation to *E. coli* culture – by V.G. Gostev, 1950. The

concentration of immunoglobulins of classes G, M, A was studied according to Mancini et al., 1965, bactericidal activity of blood serum (BABS) - by the method O.V. Smirnova and T.A. Kuzmina's method, 1966, the lysozymic activity of blood serum (LASK) - according to V.G. Dorofeychuk, 1968. The content of T-lymphocytes was determined according to M. Jondal, 1973, B-lymphocytes - according to N.S. Merdes, 1973. The incidence and safety of piglets were determined by the results of daily observations and clinical examination.

## RESULTS

The average value of the microclimate parameters in the pens where the experimental animals were kept fluctuated within the limits of: the air temperature - 20-18°C, relative humidity - 64-78%, air velocity – 0.15-0.30 m/s, concentration of harmful gases: 15-20 mg/m<sup>3</sup> for ammonia, 0.15-0.25 l/m<sup>3</sup> for carbon dioxide and 70-86 thousand CFU / m<sup>3</sup> of air for bacterial contamination.

The live weight and growth intensity of the piglets are the integral index of their natural resistance (Table 1)

Table 1 Change in live weight and live weight gains of the experimental piglets (M + m, n = 5)

Group	Age of piglets, days					SSP, g	weight gain, kg
	1	10	20	30	60		
K	1.10±0.07	2.47±0.19	4.72±0.11	7.08±0.21	15.48±0.24	239.6±10.46	14.38±0.20
0-1	1.09±0.09	2.70±0.15	5.19±0.20	7.84±0.18	17.11±0.17*	267.0±17.30	16.02±0.18
0-2	1.12±0.12	2.81±0.20	5.87±0.11	8.10±0.15*	17.86±0.22**	279.0±15.19	16.74±0.21

The studies have shown (Table 1) that the pigs from 0-2 group grew most intensively (P<0.01), the animals from group 0-1 grew less intensively, which, in our opinion, was conditioned by the manifestation of dyspepsia and bronchopneumonia. The animals from group 0-2 exceeded the piglets from the control group on the 20<sup>th</sup> day by the average daily weight gain by 24.3%, on the 30<sup>th</sup> day - by 14.4%, from group 0-1 by 9.9 and 10.7%, respectively. It should be noted that the viability of pigs from the experimental groups

was higher that can be proved by the morbidity rate: in group 0-2 it was 5%, in group 0-1-10%, in the control group -60% with disease duration from 2.0±0.1 to 3.5±0.5 and 7.2±0.5 days, respectively.

The stimulating effect of biological products on the immunological parameters of pigs has been reported [6,7,9]. We studied the dynamics of the CEC, T-and-B-lymphocytes in pigs, kept in conditions of uncontrolled microclimate (Table 2).

Table 2 Dynamics of immunological parameters of blood in piglets (M + m, n = 5)

Indicators	Group	Initial data	Research, in days				
			5	10	20	30	60
CEC, g/l	K	17.24±0.38	17.50±0.22	17.41±0.61	17.31±0.70	17.28±0.51	18.01±0.43
	0-1	17.22±0.40	17.33±0.44	17.67±0.29*	18.04±0.70*	19.84±0.42*	19.24±0.40
	0-2	17.44±0.38	17.59±0.50	18.36±0.41*	18.49±0.62*	19.48±0.33	20.76±0.43*
T-lymphocytes ml	K	4.38±0.21	4.28±0.09	3.97±0.70	4.52±0.11	4.50±0.12	4.39±0.12
	0-1	4.48±0.10	4.57±0.12	5.02±0.14*	5.17±0.12*	5.21±0.11	5.46±0.13*
	0-2	4.19±0.20	4.32±0.55	4.77±0.09*	5.22±0.61*	5.24±0.37*	5.39±0.11
B-lymphocytes ml	K	1.93±0.09	1.90±0.08	2.01±0.08	2.13±0.09	2.03±0.08	1.99±0.09
	0-1	2.01±0.09	2.16±0.03	1.93±0.01	2.30±0.03*	2.34±0.02*	2.37±0.03*
	0-2	1.95±0.04	2.02±0.07	2.08±0.06	2.16±0.01*	2.30±0.05*	2.36±0.07*

The studies have shown that the level of the CEC in the animals in the 0-1 group was higher by 4.76% - on the 10<sup>th</sup> day of the experiment, by 15, 20% on the 30<sup>th</sup> day, by 11.7% on the 60<sup>th</sup> day as compared with the control ones. In the pigs from the 0-2 group, this indicator increased by 4.95% on the 5<sup>th</sup> day, by: 6% on the 20<sup>th</sup> day, by 19.48% on the 30<sup>th</sup> day, by 11.7% on the 60<sup>th</sup> day. A significant difference was found out in the increase of T-lymphocytes in the animals from the 0-1 and - 0-2 groups from the 10<sup>th</sup> day, B-lymphocytes - on the 20<sup>th</sup> day of the experiment.

In the pigs from the 0-1 group BASK was within 56.52±2.03%, which is 3.11% lower (on the 60<sup>th</sup> day of the trial). As for the cellular indices of protection: FAN fluctuated during the whole period of the studies from 53.10±0.3 to 53.79±0.4%, the frequency of FV - from 4.28±0.09 – 4.49±0.05 (0 -1) to 4.59±0.20 – 4.93±0.04 (0-2).

Based on the results of the studies, it was found out that the preparations of RBS and Immunolac do not cause immunosuppression, they are not dangerous at the doses of 0.05 ml/mg of body weight, they improve the general state of the body and do not cause any changes indicating a pathological condition and growth depression. In the conditions of the above microclimate the adaptation process was faster in the pigs that were given the above drugs and it makes it possible to recommend them as the drugs that increase the natural resistance of the animal body.

## CONCLUSIONS

When piglets were grown at the temperature lower than 3.5-8°C, humidity - by 5-8% higher, the parameters of NH<sub>3</sub> and CO<sub>2</sub> higher by 0.8-1.2%, the contamination microflora was twice as much as recommended by the sanitary norms the genetic productive potential of the animals is not fully realized, while:

- the highest daily average weight gain was in the pigs, which were injected intramuscularly with RBS growth stimulant. They exceeded the animals from the control group on the 20<sup>th</sup> and 30<sup>th</sup> days of the experiment - by 24.3% and 14.4%, live weight - by 15.3% (P <0.05). The growth rate in the animals that were injected with Immunolac was less expressed;

- the diseases with the symptoms of dyspepsia and bronchopneumonia were registered in 5-10% of the pigs from the experimental groups that received RBS and Immunolac that is 10-12 times less as compared to the control group;

- in the pigs from the experimental group 1, the BASK was at the level of the average values of 56.52±2.03%, from the experimental group 2 – 59.48 ± 1.80%, by 5.2% higher (P<0.05);

- LASK indices in the animals of the experimental group tended to increase but the differences were not trustworthy (P<0.5);

- the cell indices of resistance in the pigs that received RBS were higher: the phagocytic activity of blood serum - by 4.3% (P<0.05), phagocytic number - by 9.7% (P<0.05);

- the highest concentration of immunoglobulin of class Jg G ( $21.82 \pm 0.3$  mg/ml) was revealed in the animals that received RBS preparations, the concentration of the above immunoglobulins was slightly below  $-19.86 \pm 0.18$  mg/ml in the pigs that received Immunolac (0-1). For immunoglobulins of class Jg M and Jg A, the fluctuations between the experimental groups were within the limits of  $2.23 \pm 0.01$  and  $2.36 \pm 0.09$  mg/ml.

bronkhopnevmoniiye, soderzhashikhysya v razlichnykh mikroklimaticheskikh usloviyakh. Problemy gigiyeny sel'skokhozyaystviykh zhivotnykh v usloviyakh intensivnogo vedeniya zhivotnovodstva: Materialy nauchno-prakticheskoy konferentsii, posvyashchenny 70-letiyu kafedry zoogigiyeny (23-24 oktyabrya 2003) Vitebsk, s 124-125.

[9]. Chumachenko, V. E., 1990. Opredeleniye yestestvennoy rezistentnosti i obmena veshestv sel'skokhozyaystviykh zhivotnykh. Kiev. 200 s.

## REFERENCES

- [1]. Adamenko, G.P., 1981. Modifikatsiya metoda Mancini dlya kolichestvennogo opredeleniya immunoglobulinov: Laboratornoye delo, № 6 s. 371-372.
- [2]. Vakar, A.N., 2004. Morfologicheskkiye i biokhimicheskkiye pokazateli krovi porosyat-sosunov pri primenenii preparatov «Dostim» i «Mastim»: sostoyaniye i problemy veterinarnoy sanitaria gigiyeny i ekologii v zhivotnovodstve: Materialy mezhdunarodnoy nauchno-prakticheskoy konferentsii. Cheboksary, s 273-274.
- [3]. Vasil'yeva, E.A., 1974. Klinicheskaya bioximiiya sel'skokhozyaystviykh zhivotnykh. M., 192 s.
- [4]. Maslyanko, R.P., I.I. Oleksyuk, & A.I. Podov'skiy, 2001. Metodichny rekomendatsyy dlya otsynky ta kontrolya immunnogo statusu tvaryn: vyznachennyya faktorov nespetsifichnoi rezistentnosti klitinnikh i gumoral'nykh mekhanizmov proti infektsiynikh zakhvoryuvan'. L'viv. 87 s.
- [5]. Masalov, V. M., O.A. Mikhaylova, & E.A. Mikheyeva, 2009. Vliyeniye vitaminnykh nastoyev na formirovaniye immuniteta porosyat. Aktual'nyye problemy veterinarnoy meditsiny: Sbornik nauchnykh trudov posvyashchennykh 90-letiyu MGAB MiB. Moskva. S 132-135.
- [6]. Ogorodnik, N.V., 2011. Vpliv preparative «Lipavit» ta «Trivit» na nespetsifichnu rezistentnost' krovi svinomatok i porosyat. Naukovo-tekhnichnyy byuleten. vipusk 12. № 3,4 - L'viv. s 288-293.
- [7]. Petryankin, F.P., 2010. Primneniye immunotropnykh preparatov dlya profilaktiki i lecheniya bolezney porosyat. Sovremennyye problemy intensivifikatsii proizvodstva svininy v stranakh SNG: Sbornik nauchnykh trudov mezhdunarodnoy nauchno prakticheskoy konferentsii po svinovodstvu (7-10 iyulya) Ul'yanovsk, T.2. S 333-337.
- [8]. Sokolov, G. A., N.M. Malinets, & I.V. Shadenko, 2003. Lecheniye telyat, bol'nykh