

EFFECT OF TILLAGE AND ZINC APPLICATION METHODS ON WEEDS AND YIELD OF MAIZE

Muhammad Azim KHAN¹, Muhammad ARIF²,
Khan Bahadar MARWAT¹

¹NWFP Agricultural University Peshawar-25130,
Pakistan, Department of Weed Science
e-mail: ahmadzaipk@yahoo.com

²NWFP Agricultural University Peshawar-25130,
Pakistan, Department of Agronomy

Field trials were conducted at Agricultural Research Farm, NWFP Agricultural University Peshawar during summer 2006 to investigate the effect of tillage and Zinc (Zn) application methods on maize yield and its associated weeds. The experiment was laid out in randomized complete block design with split-plot arrangement having three replications. The main plot consisted of conventional tillage (CT) and reduced tillage (RT) while Zn application methods were assigned to the subplots which included seed priming (dry seed, soaking seed in water, 0.01, 0.02 and 0.03% Zn solutions), foliar spray of 0.01% Zn solution, soil application at the rate of 5 kg ha⁻¹ and combination of soil application (@ 5 kg ha⁻¹) plus foliar spray (0.01% Zn solution). Tillage significantly affected weed density (WD) and yield of maize. CT suppressed WD recorded 21 and 42 days after sowing (DAS). Likewise, CT also suppressed the major weeds (Cyperus rotundus, Digitaria sanguinalis, and Convolvulus arvensis). Higher grain yield was recorded in CT plots as compared to RT. Zinc application methods significantly affected WD recorded 42 days after sowing and grain yield of maize. The interaction between tillage and Zn application methods for WD recorded 42 DAS was also significant. Greater WD was recorded in RT as compared to CT. Mean of the interaction values indicated that WD was lower in CT as compared to RT irrespective of Zn application methods. Yield was higher in CT as compared to RT. Likewise water soaking and Zn application methods improved maize yield as compared to control. It is concluded that CT resulted in lower WD and higher yield of maize. Furthermore water soaking and Zn application methods enhanced grain yield of maize as compared to control.

Keywords: soil tillage, weed, maize, zinc.

Tillage is considered the most effective farm activity for the purpose of developing a desired soil structure. It improves the physical conditions of soil and favors the rooting characteristics of plants, which lead to an enhanced nutrient uptake and better yield of crops. Weeds are probably the most ever-present class of crop pests and on the odd occasion cause massive crop failures over vast areas. They reduce the crop yield and deteriorate the quality of produce and hence reduce

the market value of the turn out (Arif *et al.*, 2006). They use the soil fertility, available moisture, nutrient and compete for space and light with crop plant, which result in yield reduction (Khan *et al.*, 2002 and 2004). If left uncontrolled, the weeds in many fields are capable of reducing yields by more than 80% (Karlen *et al.*, 2002).

The composition of weed communities is greatly affected by tillage but it is difficult to control weeds in reduced tillage (RT). RT often favors annual grasses and discourages annual dicotyledonous species (Froud-Williams *et al.*, 1981; Gill and Arshad, 1995). However, generalizations are limited, because the effect of tillage on annual weeds is species-specific (Buhler, 1992), and the same species may respond differently when soil properties and other site characteristics vary. A study carried out in Saskatchewan, Canada, revealed that the influence of location and year on changes in weed communities was more as compared to tillage systems (Derksen *et al.*, 1993). Increased soil disturbance decreased the number of weed species and species diversity in maize cropping (Cardina *et al.*, 1991). The relative contributions to the size and diversity of weed flora were greater by common species under conventional tillage (CT) and by rare species in less intensive tillage systems in spring crops (Gill and Arshad, 1995).

The disadvantages of RT are infestations by several annual and perennial species and rapid increase of the seed bank near the soil surface. Hence, occasional or rotational use of RT may be a practical way to adopt RT into CT systems. Changes between tillage practices, from CT to RT and vice versa, were effective in suppressing weed growth and preventing seed accumulation (Nakamoto *et al.*, 2006).

The response of crop yield to tillage practices is rather contradictory. Zuegec (2003) reported that the highest and the most stable maize grain yields were obtained with CT. Whereas Hussain *et al.* (1999) reported that four year average maize yields were equal for no tillage, chisel plow and moldboard plow systems and four year average soybean yield with no tillage was 15% higher than with the moldboard plow systems. Similarly, Tarkalson *et al.* (2006) reported that no-till system increased long term average winter wheat and grain sorghum yields compared to CT.

There are two reasons to increase the micronutrient content of the grains of major food crops: enhancement of the agronomic productivity of the crop and improvement of the nutritional value of staple foods for humans. It has been estimated that roughly 40 % of people throughout the world suffer from micronutrient malnutrition, most commonly from iron, zinc (Zn), iodine, or various vitamin deficiencies (Welch *et al.*, 1997).

There are three main methods of applying micronutrients to crops: soil fertilization, foliar sprays and seed treatment. Foliar applications of micronutrient sprays are effective towards both goals (Wilhelm *et al.*, 1988; Savithri *et al.*, 1999), but this method is too costly to be widely practiced by resource-poor farmers in some regions because of the amount of fertilizer, equipment and labour required for repeated spraying. Likewise, the difficulty in obtaining high quality

micronutrient fertilizers and spreading them evenly on the soil can be unaffordable. Treating seeds with micronutrients potentially provides a simple inexpensive method for improving micronutrient plant nutrition. Farmers in south Asia have responded favorably to seed priming, a simple technology of soaking seeds overnight in water prior to sowing (Harris, 1996; Harris *et al.*, 1999). Seed priming in water has been shown to decrease time between sowing and emergence and to improve seedling vigour (Harris, 1996; Parera and Cantliffe, 1994). Priming seeds in solutions of macro or micronutrients has been shown to improve yield of rice (Peeran and Natanasabapathy, 1980), wheat (Khalid and Malik, 1982; Marcar and Graham, 1986; Wilhelm *et al.*, 1988) and forage legumes (Sherrell, 1984), but the potential to damage the seed and inhibit germination by priming at high nutrient concentrations has also been reported (Roberts, 1948).

The instant project was initiated with the objectives to study the effect of tillage and Zn application methods on weeds and grain yield of maize.

MATERIALS AND METHODS

Experimental Site

Field experiment was conducted at Agricultural Research Farm, NWFP Agricultural University, Peshawar, Pakistan during summer 2006. The experimental site is located at 34° N, 71.3° E and an altitude of 347 m ASML. The soil of the field was clay loam, low in nitrogen (0.03-0.04%), low in organic matter (0.7-0.9%) and alkaline in reaction (pH 8.0-8.2). The average monthly temperature, relative humidity, rainfall and soil temperature data during the growing season of the crop are reported in Fig 1.




Fig 1. Average monthly temperature, relative humidity, rainfall and soil temperature for May to August 2005. Source: Weather Station, NWFP Agricultural University, Peshawar, Pakistan.

Experimental procedure

The experiment was laid out in Randomized Complete Block design with split plot arrangement and three replications. The main plot consisted of conventional tillage (CT) and reduced tillage (RT). Zn application methods were assigned to the subplots which included seed priming (dry seed, soaking seed in water, 0.01, 0.02 and 0.03% Zn solutions), foliar spray of 0.01% Zn solution at 5-6 leaf stage, soil application at the rate of 5 kg ha⁻¹ before sowing and combination of soil application at the rate of 5 kg ha⁻¹ before sowing plus foliar spray of 0.01% Zn solution at 5-6 leaf stage. Dry seed of maize was used as control treatment. Zinc sulphate (ZnSO₄·7H₂O) was used as source of Zn. Nitrogen and phosphorus were applied at the rate of 120 and 100 kg ha⁻¹. Urea and single super phosphate were used as sources of N and P, respectively. Certified category seed of maize variety 'Azam' was procured from Cereals Crops Research Institute (CCRI), Nowshera, NWFP and was sown with the help of a hand hoe in 15 m² plot at the rate of 60 kg ha⁻¹. The row to row distance was 70 cm while plant to plant distance of 20 cm was maintained. All other agronomic practices were carried out uniformly in all plots. The data on weed density (WD) were recorded 21 and 42 days after sowing (DAS).

Statistical Analysis

The data were statistically analyzed using ANOVA and means were separated using LSD test at 5% level of probability (Steel and Torrie, 1984).

RESULTS AND DISCUSSION

Weed density m⁻² (21 DAS)

The effect of tillage on weed density was significant whereas the effect of Zn application methods was not significant. The interaction between the two factors for WD 21 DAS was also not significant (Table 1). Weed density was higher in RT as compared to CT (Fig. 1). The higher weed density in RT was obviously because of seed bank which was not disturbed to the greater extent. Moreover most of the weed seeds (71%) are concentrated at a depth of 10-15 cm therefore deep and repeated tillage affected the weed seed bank (Nakamoto *et al.*, 2006; Swanton *et al.*, 2000). The results are in line with Cardina *et al.* (1991) who reported that density of weed populations may increase under RT. These results also confirm the findings of Khattak and Khan (2005) who stated that with increasing the number of tillage, weed density was decreased due to destruction of the germinated annual and perennial weeds.

Table 1
Mean squares for weed densities and grain yield as affected by tillage and Zn application methods

S.o.V.	df	Weed Density 21 DAS	Weed Density 42 DAS	Grain yield	<i>Convolvulus arvensis</i>	<i>Cyperus rotundus</i>	<i>Digitaria sanguinalis</i>
Mean Squares							
Replication	2	67.69	1667.2	97258	4.88	141.6	77.0
Tillage	1	19240*	83792.3*	1269179*	584.5*	59361.3*	3176.9*
Error I	2	40.77	799.3	40946	0.318	42.3	89.3
Zinc	7	75.64	1253.6*	1747029*	15.94	1472.3	78.8
T x Zn	7	127.21	2967.6*	252920	10.16	1479.3	73.9
Error II	28	147.87	420.2	305516	9.77	627.2	35.4

* Significant at 5% level of probability

Weed density m⁻² (42 DAS)

Perusal of the data revealed that the effects of tillage, Zn application methods and interaction between the two factors were significant (Table 1). Greater weed density was recorded in RT as compared to CT (Fig. 1). It is obvious from the data that WD 42 DAS was higher than WD recorded 21 DAS. It can be implied that all weeds did not germinate till 21 DAS and weeds germinated with the passage of time. Major weeds infested the experimental field were *Cyperus rotundus*, *Convolvulus arvensis* and *Digitaria sanguinalis*. These results suggested that perennial as well as annual weeds can be suppressed by tillage. The disadvantages of RT are infestations by several annual and perennial species and rapid increase of the seed bank near the soil surface. These disadvantages became more prevalent with time and may hinder the continuous long term use of RT without herbicides applications. Analogous results were reported by Nakmoto *et al.* (2006) who reported that switching from CT to RT and vice versa, were effective in suppressing weed growth and preventing seed accumulation. Maximum WD (140 m⁻²) was recorded for foliar spray of Zn while minimum WD (95 m⁻²) was noted for seed priming in 0.02% Zn solution. Mean values of the interaction indicated that WD was lower in CT as compared to RT irrespective of Zn application methods (Table 2).

Table 2

Weed density (42 DAS) as affected by tillage systems and Zn application methods

Zn application methods	Reduced Tillage	Conventional Tillage	Mean
Tillage x Zn application methods			
Dry seed	124 de	84 fg	104 bc
Water soaking	157 bcd	70 fgh	113 bc
0.01% Zn priming	161 bc	85 fg	123 ab
0.02% Zn priming	138 cd	51 gh	95 c
0.03% Zn priming	182 b	43 h	112 bc
Foliar Spray	218 a	62 fgh	140 a
Soil application	123 de	73 fgh	98 c
Foliar Spray + Soil application	124 de	90 ef	107 bc
LSD	34.28		24.24

Means in a column followed by the same letters are not significantly different at alpha 0.05 using LSD test.

Density of major weeds

Density of major weeds 42 DAS was significantly affected by tillage (Table 1). However, the effect of Zn application methods was not significant. Likewise the interaction between the two factors for density of major weeds was also not significant. Major weeds included *Cyperus rotundus*, *Digitaria sanguinalis* and *Convolvulus arvensis*. CT suppressed densities of all major weeds (Fig. 2). The results are in line with those of De La Fuente *et al.* (1999) who reported that tillage was a major variable influencing weed communities in soybean fields. Similar results were reported by Staniforth and Wiese (1985); Frick and Thomas (1992) who found annual and perennial weed at lower densities in CT than in RT. Many

researchers like Froud-Williams *et al.* (1981) are of the opinion that introduction of new tillage practices (reduced, minimum or non-tillage) commonly causes changes in the composition and abundance of weed species present in cropping systems. The results are in agreement with Swanton *et al.* (1999) who reported that disturbance caused by tillage was more important than nitrogen rate and cover crop as a mechanism influencing the composition of the weed flora within mono-cropped maize. However, according to Pollard *et al.* (1982) the response of weeds to tillage is variable. Likewise, the results are not agreed by Derksen *et al.* (1993) who suggested that changes in weed communities were influenced more by environmental factors (location and year) than by the tillage systems.




Fig 1. Effect of tillage on weed density recorded 21 and 42 DAS. Vertical bars denote Standard Error




Fig 2. Effect of tillage on density of *Cyperus rotundus*, *Digitaria sanguinalis* and *Convolvulus arvensis*

Grain yield (kg ha⁻¹)

Tillage and Zn application methods had significant effect on grain yield of maize, whereas their interaction was not significant (Table 1). Yield was higher in CT compared to RT (Table 3). This phenomenon may be attributed to difference in WD between the tillage treatments. Greater WD occurred in the RT plots and that may have influenced maize yield. Similar results were reported by Khattak *et al.* (2005) who found that deep tillage significantly decreased weed density and increased the crop yield. These results do agree with the findings of Zugec (1986)

who obtained the highest and the most stable grain yields with conventional tillage. However the results are not in line with Hussain *et al.* (1999) who concluded that crop yields for eight years (four years maize and four year soybean) appeared to show improved long term productivity of no-till compared with that of moldboard plow and chisel plow systems. Tarkalson *et al.* (2006) are also not supportive of these findings that no-till increased long term average winter wheat and grain sorghum yields compared with conventional tillage.

Water priming produced higher grain yield (37%) as compared to control. The yield increment due water priming has been reported by many researchers who concluded that 'on-farm' seed priming with water alone is effective in substantially increasing yields of chickpea (Harris *et al.*, 1999; Musa *et al.* 2001), wheat and maize (Harris *et al.*, 2001). Seed priming in 0.01, 0.02 and 0.03% Zn solutions produced 33, 34 and 36% more grain yield than control. Foliar spray of 0.01% Zn solution and soil application at the rate of 5 kg ha⁻¹ produced 30 and 41% greater grain yield than control. Soil application (at the rate of 5 kg ha⁻¹) along with foliar spray (at the rate of 0.01% Zn solution) resulted in the maximum grain yield which was 44% higher than the control.

Soil applied Zn resulted in higher grain yield of maize as compared to seed priming and foliar spray. Similar results were reported by Yilmaz *et al.* (1997) who concluded that soil applied Zn was a superior fertilization method compared with Zn treated wheat seed or foliar Zn applications. Similar results were reported by Shah *et al.* (1985) and Rehman and Barnard (1988) who obtained maximum yields in maize and lentil with the application of 10 and 5.5 kg Zn ha⁻¹ as compared to control, respectively. Likewise application of foliar-sprayed Zn (50 ppm) significantly increased cotton yield as compared to untreated plants (Zeng QingFang., 1996). The results obtained are also confirmed by Sawan *et al.* (1988).

Table 3

Effect of tillage and Zn application methods on yield (kg ha⁻¹) of maize

Zn application methods	Reduced tillage	Conventional tillage	Mean	% increase over control
	Tillage x Zn application methods			
Dry seed	2619	3113	2866 b	
Water	3214	4246	3730 ab	37
0.01% Zn priming	3433	3204	3318 ab	33
0.02% Zn priming	3108	3632	3370 ab	34
0.03% Zn priming	3460	3819	3639 ab	36
Foliar Spray	3019	2912	2965 ab	30
Soil application	4095	4127	4111 ab	41
Foliar Spray + Soil application	4188	4687	4437 a	44
Mean	3392	3717		
LSD			1563	

Means in a column followed by the same letters are not significantly different at alpha 0.05 using LSD test.

REFERENCES

- Arif, M., M.A. Khan, H. Akbar, Sajjad and S. Ali. 2006. Prospects of wheat as a dual purpose crop and its impact on weeds. *Pakistan Journal of Weed Science Research*, 12(1-2):13-17.
- Buhler, C.D. 1992. Population dynamics and control of annual weeds in corn (*Zea mays*) as influenced by tillage systems. *Weed Science*, 40:241-248.
- Cardina, J., E. Regnier, K. Harrison. 1991. Long-term tillage effects on seed banks in three Ohio soils. *Weed Science*, 39:186-194.
- De La Fuente, E.B., S.A. Suarez, C.M. Ghera and R.J.C. Leon. 1999. Soybean weed communities: relationship with cultural history and corn yield. *Agronomy Journal*, 91:409-417.
- Derkson, D.A., G.P. Lafond, H.A. Loeppky and C.J. Swanton. 1993. impact of agronomic practices on weed communities: Fallow within tillage systems. *Weed Science*, 42:729-735.
- Frick, B. and A.G. Thomas. 1992. Weed surveys in different tillage systems in southwestern Ontario fields crops. *Canadian Journal of Plant Science*, 72:1337-1347.
- Froud-Williams, R.J., R.J. Chancellor and D.S.H. Drennan. 1981. Potential changes in weed flora associated with reduced-cultivation systems in cereal production in temperate regions. *Weed Research*, 21: 99-109.
- Gill, K.S. and M.A. Arshad. 1995. Weed flora in the early growth period of spring crops under conventional, reduced and zero tillage systems on a clay soil in northern Alberta, Canada. *Soil and Tillage Research*, 33:65-79.
- Harris, D. 1996. The effects of manure, genotype, seed priming, depth and date of sowing on the emergence and early growth of *Sorghum bicolor* (L.) Moench in semi-arid Botswana. *Soil and Tillage Research*, 40:73-88.
- Harris, D., A. Joshi, P.A. Khan, P. Gothkar and P.S. Sodhi. 1999. On farm seed priming in semi arid agriculture: Development and evaluation in maize, rice and chickpea in India using participatory methods. *Experimental Agriculture*, 35:15-29.
- Hussain, I., K.R. Olson, and S.A. Ebelhar. 1999. Impacts of tillage and no-till on production of maize and soybean on an eroded Illinois silt loam soil. *Soil and Tillage Research*, 52:37-49.
- Karlen, L.D., D.D. Buhler, M.M. Ellusbury and S.S. Andrews. 2002. Soil, weeds and insect management strategies for sustainable agriculture. *Journal of biological Sciences*, 2(1):58-62.
- Khan, M.A., K.B. Marwat, G. Hassan and W.A. Shah. 2002. Effect of different weed free periods on the growth and yield of wheat. *Pak. J. Agri., Agril. Eng. Vet. Sci.* 18(1-2):30-33.
- Khan, I., G. Hassan, M.I. Khan and I.A. Khan. 2004. Efficacy of some new herbicidal molecules on grassy and broadleaf weeds in wheat-II. *Pakistan Journal of Weed Science Research*, 10(1-2):33-38.
- Khalid, B. Y. and Malik, N.S.A. 1982. Prsowing soaking of wheat seeds in copper and manganese, solutions. *Journal of Communications in Soil Science and Plant Analysis*. 13:981-986.
- Khattak, M.K., M.J. Khan, M. Asrar, S. Wahab and M. Ramzan. 2005. Enhancement of wheat yield and yield components by controlling weeds with deep tillage practices under irrigated clay loam soil conditions. *Pakistan Journal of Weed Science Research*, 11: 131-136.
- Khattak, M.K. and M.J. Khan. 2005. Effect of different tillage practices on weeds and yield of chickpea under sandy loam soil conditions. *Pakistan Journal of Weed Science Research*, 11: 157-164.
- Marcar, N.E. and Graham, R.D. 1986. Effect of seed manganese content on the growth of wheat (*Triticum aestivum*) under manganese deficiency. *Plant and Soil*. 96:165-173.

- Musa, A.M., D.Harris, C. Johansen, and J. Kumar. 2001. Short duration chickpea to replace fallow after aman rice: the role of on-farm seed priming in the High Barind Tract of *Bangladesh Journal of Experimental Agriculture*, 37(4): 509-521.
- Nakamoto, T., J. Yamagishi, F. Miura. 2006. Effect of reduced tillage on weeds and soil organisms in winter wheat in summer maize cropping on humic andosols in central Japan. *Soil and Tillage Research*, 85:94-106.
- Parera, C.A. and D.J. Cantliffe. 1994. Pre-sowing seed priming. *Horticultural review*, 16:109-141.
- Peeran, S.N. and S. Natanasabapathy. 1980. Potassium chloride pretreatment on rice seeds. *International Rice Research Newsletter*. 5:19.
- Pollard, F., S.R. Moss, G.W. Cussans and R.J. Froud-Williams. 1982. The influence of tillage on weed flora in a succession of winter wheat crops on a clay loam soil and silt loam soil. *Weed Res.* 22:129-136.
- Rehman, G. and E.E. Barnard. 1988. Yield, quality and agronomic characteristics of lentil with phosphorus and zinc. *Sarhad Journal of Agriculture*, 4:233-240.
- Roberts, W.O. 1948. Prevention of mineral deficiency by soaking seed in nutrient solution. *Journal of Agricultural Sciences*, 38:458-468.
- Savithri, P., R. Perumal and R. Nagarajan. 1999. Soil and crop management technologies for enhancing rice production under micronutrient constraints. *Nutrient Cycling in Agroecosystems*. 53:83-92.
- Shah, M., J.K. Khattak and S.A. Rizvi. 1985. Effect of boron and zinc fertilization on the yield of maize. *Sarhad Journal of Agriculture*, 1:205-210.
- Sherrell, C.G. 1984. Effect of molybdenum concentration in the seed on the response of pasture legumes to molybdenum. *Newzealand Journal of Agricultural Research*, 27:417-423.
- Staniforth, D.W. and A.F. Wiese. 1985. Weed biology and its relationship to weed control in limited-tillage systems in: *Weed control in Limited Tillage Systems* (ed. AF Wiese), 15-25. Monograph Series of the Weed Science Society of America, Champaign, IL, USA.
- Steel, R.G.D. and J.H. Torrie. 1984. Principles and procedures of statistics, 2nd ed., p.172-177. McGraw Hill Book Co., Singapore.
- Sawan, Z. M., A. A. El-Farra and Z. A. Mohamed. 1988. Effect of nitrogen fertilization, foliar application of calcium and some micro-elements on cottonseed, protein and oil yields and oil properties of Egyptian cotton. *Annalo Botanica* (Rome), 46:167-174.
- Swanton, C.J., A. Shrestha, S.Z. Knezevic, R.C. Roy and B.R. Ball-Coelho. 1999. Effect of tillage systems, N and cover crop on the composition of weed flora. *Weed Science*, 47:454-461.
- Swanton, C.J., A.Shrestha, S.Z. Knezevic, R.C. Roy and B.R. Ball-Coelho. 2000. Influence of tillage type on vertical weed seed bank distribution in a sandy soil. *Canadian Journal of Plant Science*, 80:455-457.
- Tarkalson, D.D., G.W. Hergert and K.G. Cassman. 2006. Long-term effects of tillage on soil chemical properties and grain yields of a dryland winter wheat-sorghum/corn-fallow rotation in the great plains. *Agronomy Journal*, 98:26-33.
- Welch, R.M., G.F. Combs (Jr) and J.M. Duxbury. 1997. Toward a "greener" revolution. *Issues in Science and Technology*, 14:50-58.
- Wilhelm, N.S., R.D. Graham and A.D. Rovira. 1988. Application of different sources of manganese sulphate decreases take-all (*Gaeumannomyces graminis* var. *tritici*) of wheat grown in a manganese deficient soil. *Australian Journal of Agricultural Research*, 39:1-10.
- QingFang, Z.1996. Research on the effect of zinc applied to calcareous soil in cotton. *China Cottons*. 23:21.
- Yilmaz, A., H. Ekiz, I. Gultekin, B. Torun, H.K. Barut, and I. Cakamakl. 1997. Effect of seed zinc content on grain yield and zinc concentration of wheat grown in zinc-deficient calcareous soils. *Journal of Plant Nutrition*, 21:2257-2264.
- Zugec, I. 2003. The effect of reduced soil tillage on maize (*Zea mays* L.) grain yield in Eastern Croatia (Yugoslavia). *Soil and Tillage Research*, 7:19-28.