

RESEARCHES REGARDING THE ENTOMOFAUNA OF COLEOPTERAS WITHIN SOME AGRICULTURAL CROPS FROM THE NORTH PART OF PORTUGAL

Valentin Teodor TUDORACHE¹, Mihai TĂLMACIU¹, Nela TĂLMACIU¹, Monica HEREÀ¹

e-mail: tudorachevalentin2007@gmail.com

Abstract

The purpose of this research was that to identify the useful entomofauna of Coleopteras from the agricultural crops which have been taken into the study: olive trees and orange trees. The researches were carried out within the Quinta da Pinhao station, which belongs to the University of "Tras os Montes e Alto Douro". The biological material was sampled by the mean of D-Vacuum at 20th of June of 2017, from each 6 trees belonging to each crop, which have been aimlessly chosen. The D-Vacuum has worked one minute for each tree taken into the study. After carrying out the experiments, all the sampled insects were brought into the Laboratory of Entomology in order to be counted and determined.

From the *Coleoptera* order, species with the biggest number of samples were:

- within the olive trees: *Chrysomelidae* family-91 samples, *Coccinellidae*- 56 samples, *Aderidae*-42 samples and *Scarabaeidae*-38 samples. From the *Coccinellidae* family, species with the biggest number of samples were: *Coccinella septempunctata* (31) and *Hippodamia variegata* (16).
- within the orange trees: *Chrysomelidae* family-142 samples and *Coccinellidae* family- 54 samples.. From the *Coccinellidae* family, species with the biggest number of samples was: *Coccinella septempunctata* (33) .

Key words: D-Vacuum, *Chrysomelidae*, *Scarabaeidae*, *Coccinellidae*.

The insects of *Chrysomelidae* family are commonly known as leaf beetles and include more than 50.000 species, making up one of the largest and most common encountered of all beetle families. Leaf beetles are partially recognizable by their tarsal formula, which appears to be 4-4-4, but is actually 5-5-5 as the fourth tarsal segment is very small and hidden by the third one. Adult and larval leaf beetles feed on all sorts of plant tissue, and all species are fully herbivorous. Many are serious pests of cultivated plants (eg. *Leptinotarsa decemlineata*-Colorado potato beetle).

Coccinellidae family has species of round or oval species, almost hemispherical, with spots on the wings with color spots and contrasting patterns. The most species of *Coccinellidae* are beneficial predators which prefer the aphids as main feed.

According to the author (Foltz, 2002), there are more than 5.000 species all around the world. The most representative species of the *Coccinellidae* family are: *Coccinella septempunctata*, *Hippodamia variegata*, *Propylaea quatuodecimpunctata*.

MATERIAL AND METHOD

In order to carry out the researches, the insects were sampled from one station: Quinta da Pinhao, belonging to University of "Tras os Montes e Alto Douro", by using the method of D-Vacuum. The experiments were carried out at 20th of June of 2017. Each 6 trees belonging to olive and orange orchards were taken into this study, the vacuum working 1 minute for each tree. After finishing the experiments, all the samples were brought into the laboratory, counted and determined by the mean of binocular magnifier.

RESULTS AND DISCUSSIONS

The samples of the biological material were carried out at 20th of June of 2017. Within the olive trees orchard, the situation regarding the entomofauna of Coleopteras looks as follows (*table 1*):

-Tree number 1: there were 83 samples identified from Coleoptera order, belonging to *Chrysomelidae* family (22), *Coccinellidae* family (21), *Scarabaeidae* (15), *Staphylinidae* (11),

¹"Ion Ionescu de la Brad" University of Agricultural Sciences and Veterinary Medicine, Iași

Aderidae (14). From the *Coccinellidae* family, 13 species, 5 belong to *Hippodamia variegata* species, and 3 samples belong to *Propylaea quatordecimpunctata* species;

-Tree number 2: there were 51 samples identified from Coleoptera order, belonging to *Chrysomelidae* family (19), *Scarabaeidae* (8), *Mordellidae* (11) and *Coccinellidae* family (13). From the *Coccinellidae* family, 7 samples belongs to *Coccinella septempunctata* species, 4 samples belong to *Nephus quadrimaculatus* species and 2 samples belong to *Propylaea quatordecimpunctata* species;

-Tree number 3: there were 53 samples identified from Coleoptera order, belonging to *Chrysomelidae* family (20), *Aderidae* (17), *Cerambycidae* (12) and *Staphylinidae* (2) and *Coccinellidae* family (2). From the *Coccinellidae* family, the both samples belong to *Coccinella septempunctata* species;

- Tree number 4: there were 50 samples identified from Coleoptera order, belonging to

samples belong to *Coccinella septempunctata* *Chrysomelidae* family (14), *Scarabaeidae* (12), *Aderidae* (8), *Mordellidae* (7), *Coccinellidae* (6 samples) and *Cetoniidae* (3). From the *Coccinellidae* family, all the collects samples belong to *Hippodamia variegata* species;

-Tree number 5: there were 30 samples identified from Coleoptera order, belonging to *Coccinellidae* family (9), *Anthicidae* family (8), *Chrysomelidae* (4), *Aderidae* (3), *Scarabaeidae* (3); *Dermestidae* (3). From the *Coccinellidae* family, 7 samples belong to *Coccinella septempunctata* and two samples belong to *Hippodamia variegata* species;

- Tree number 6: there were 21 samples identified from Coleoptera order , belonging to *Chrysomelidae* family (12), *Cerambycidae* family (4) and *Coccinellidae* (5). From the *Coccinellidae* family, 3 samples belong to *Hippodamia variegata* species and two samples belong to *Coccinella septempunctata* species.

Table 1

Entomofauna of coleopteras sampled from olive trees by the mean of D-Vacuum

TREE	FAMILY	SPECIES	TOTAL SPECIES	TOTAL
1.	<i>Chrysomelidae</i>	-	-	22
	<i>Coccinellidae</i>	<i>Coccinella septempunctata</i>	13	21
		<i>Hippodamia variegata</i>	5	
		<i>Propylaea quatordecimpunctata</i>	3	
	<i>Scarabaeidae</i>			15
	<i>Aderidae</i>			14
	<i>Staphylinidae</i>			11
2.	TOTAL			83
	<i>Chrysomelidae</i>	-		19
	<i>Coccinellidae</i>	<i>Coccinella septempunctata</i>	7	13
		<i>Nephus quadrimaculatus</i>	4	
		<i>Propylaea quatordecimpunctata</i>	2	
	<i>Mordellidae</i>			11
	<i>Scarabaeidae</i>			8
3.	TOTAL			51
	<i>Chrysomelidae</i>			20
	<i>Aderidae</i>			17
	<i>Cerambycidae</i>			12
	<i>Staphylinidae</i>			2
	<i>Coccinellidae</i>	<i>Coccinella septempunctata</i>	2	2
	TOTAL			53
4.	<i>Chrysomelidae</i>			14
	<i>Scarabaeidae</i>			12
	<i>Aderidae</i>			8
	<i>Mordellidae</i>			7
	<i>Coccinellidae</i>	<i>Hippodamia variegata</i>	6	6
	<i>Cetoniidae</i>			3
	TOTAL			50

5.	<i>Coccinellidae</i>	<i>Coccinella septempunctata</i>	7	9
		<i>Hippodamia variegata</i>	2	
	<i>Anthicidae</i>			8
	<i>Chrysomelidae</i>			4
	<i>Aderidae</i>			3
	<i>Dermestidae</i>			3
	<i>Scarabaeidae</i>			3
	TOTAL			30
6.	<i>Chrysomelidae</i>			12
	<i>Coccinellidae</i>	<i>Hippodamia variegata</i>	3	5
		<i>Coccinella septempunctata</i>	2	
	<i>Cerambycidae</i>			4
	TOTAL			21
	TOTAL ENTOMOFAUNA OF COLEOPTERAS			288

Within the orange trees orchard, the situation regarding the entomofauna of Coleopteras looks as follows (table 2):

-Tree number 1: there were 42 samples identified from *Coleoptera* order, belonging to *Chrysomelidae* family (30), *Coccinellidae* (7) and *Curculionidae* (5). From the *Coccinellidae* family, 5 samples belong to *Coccinella septempunctata* species and two samples belong to *Coccinella 5-punctata* species;

-Tree number 2: there were 37 samples identified from *Coleoptera* order, belonging to *Chrysomelidae* family (22), *Coccinellidae* (10), *Curculionidae* (3) and *Staphylinidae* (2). From the *Coccinellidae* family, 5 samples belong to *Coccinella septempunctata* species, 3 samples belong to *Coccinella subpunctata* species and 2 samples belong to *Propylaea quatordecimpunctata* species;

-Tree number 3: there were 28 samples identified from *Coleoptera* order, belonging to *Chrysomelidae* family (16), *Cerambycidae* (4), *Scarabaeidae* (4), *Anthicidae* (2) and *Curculionidae* (2);

-Tree number 4: there were 41 samples identified from *Coleoptera* order, belonging to *hrysomelidae* family (27), *Coccinellidae* family (9), *Scarabaeidae* (3) and *Staphylinidae* (2). From the *Coccinellidae* family, 6 samples belong to *Coccinella septempunctata* species, 2 samples belong to *Coccinella subpunctata* and 1 samples belongs to *Adalia bipunctata* species;

-Tree number 5: there were 53 samples identified from *Coleoptera* order, belonging to *Chrysomelidae* family (26), *Coccinellidae* (16), *Cerambycidae* (10) and *Staphylinidae* (1). From the *Coccinellidae* family, 7 samples belong to *Coccinella septempunctata* species, 4 samples belong to *Coccinella 5-punctata*, 3 samples belong to *Adalia bipunctata* and 2 samples belong to *Coccinella subpunctata*;

- Tree number 6: there were 38 samples identified from *Coleoptera* order, belonging to *Chrysomelidae* family (21), *Coccinellidae* family (12), *Scarabaeidae* (4) and *Curculionidae* (1). From the *Coccinellidae* family, 10 samples belong to *Coccinella septempunctata* species and 2 samples belong to *Propylaea quatordecimpunctata* species.

Table 2

Entomofauna of coleopteras sampled from orange trees by the mean of D-Vacuum

TREE	FAMILY	SPECIES	TOTAL SPECIES	TOTAL
1.	<i>Chrysomelidae</i>	-	-	30
	<i>Coccinellidae</i>	<i>Coccinella septempunctata</i>	5	7
		<i>Coccinella 5-punctata</i>	2	
	<i>Curculionidae</i>			5
	TOTAL			42
	<i>Chrysomelidae</i>	-		22
2.	<i>Coccinellidae</i>	<i>Coccinella septempunctata</i>	5	10
		<i>Coccinella subpunctata</i>	3	
		<i>Propylaea quatordecimpunctata</i>	2	
	<i>Curculionidae</i>			3
	<i>Staphylinidae</i>			2
	TOTAL			37

	<i>Chrysomelidae</i>			16
	<i>Scarabaeidae</i>			4
	<i>Cerambycidae</i>			4
	<i>Anthicidae</i>			2
	<i>Curculionidae</i>			2
	TOTAL			28
3.	<i>Chrysomelidae</i>			27
		<i>Coccinella septempunctata</i>	6	
		<i>Coccinella subpunctata</i>	2	9
		<i>Adalia bipunctata</i>	1	
	<i>Scarabaeidae</i>			3
	<i>Staphylinidae</i>			2
	TOTAL			41
4.	<i>Chrysomelidae</i>			26
		<i>Coccinella septempunctata</i>	7	
		<i>Coccinella 5-punctata</i>	4	16
		<i>Adalia bipunctata</i>	3	
	<i>Cerambycidae</i>			10
	<i>Staphylinidae</i>			1
	TOTAL			53
5.	<i>Chrysomelidae</i>			21
		<i>Coccinella septempunctata</i>	10	
		<i>Propylaea quatordecimpunctata</i>	2	12
	<i>Scarabaeidae</i>			4
	<i>Curculionidae</i>			1
	TOTAL			38
	TOTAL ENTOMOFAUNA OF COLEOPTERAS			239

Table 3

Entomofauna of Coleoptera-Coccinellidae sampled within both crops (olive and oranges)

Current number	Tree	Species	No.of samples	Total
1.	olive	<i>Coccinella septempunctata</i>	31	56
		<i>Hippodamia variegata</i>	16	
		<i>Propylaea quatordecimpunctata</i>	5	
		<i>Nephus quadrimaculatus</i>	4	
2.	orange	<i>Coccinella septempunctata</i>	33	54
		<i>Coccinella subpunctata</i>	7	
		<i>Coccinella var.6-punctata</i>	6	
		<i>Propylaea quatordecimpunctata</i>	4	
		<i>Adalia bipunctata</i>	4	

Entomofauna of Coleoptera-Coccinellidae sampled within olive trees and orange trees is presented in the table 3.

During the day of 20th of June of 2017, by the mean of D-Vaccum method, species from Coccinellidae family (Coleoptera-Coccinellidae) have been sampled within Quinta da Pinhao station. Results look thus:

Within the olive orchard, 56 samples belonging to Coccinellidae family (Coleoptera-Coccinellidae) have been sampled. Species with

the biggest number of samples were: *Coccinella septempunctata* (31) and *Hippodamia variegata* (16). In total, 4 species of Coccinellidae (Coleoptera-Coccinellidae) have been identified.

Within the orange orchard, 54 samples which belong to Coccinellidae family (Coleoptera-Coccinellidae) have been sampled. Species with the biggest number of samples were: *Coccinella septempunctata* (33) and *Coccinella subpunctata* (7).

In total, 5 species that belong to *Coccinellidae* family (*Coleoptera-Coccinellidae*) have been identified. species.

CONCLUSIONS

It can be concluded that, during the day of 20th of June 2017, within the olive orchard, from a total of 288 samples of *Coleopteras*, the most significant number belongs to *Chrysomelidae* family (91), followed by *Coccinellidae* family (56), *Aderidae* (42) and *Scarabaeidae* (38), while the leastest number of samples belongs to: *Anthicidae* family (8), *Cetoniidae* and *Dermestidae* families (each 3). The most dominant species of *Coccinellidae* family are: *Coccinella septempunctata* (31 samples), *Hippodamia variegata* (16 samples), while the leastest number of samples belongs to the following species: *Propylaea quatordecimpunctata* (5 samples) and *Nephus quadrimaculatus* (4 samples).

Within the orange orchard, from a total of 239 samples of *Coleopteras*, the most significant number belongs to *Chrysomelidae* family (142), followed by *Coccinellidae* family (54), while the leastest number of samples belongs to: *Staphylinidae* family (5) and *Anthicidae* family (2). The most dominant species of *Coccinellidae* family is: *Coccinella septempunctata* species (33 samples), while the species with the leastest number of samples are: *Propylaea quatordecimpunctata* and *Adalia bipunctata* (each 4 samples).

The biggest number of samples from *Coleoptera* order was identified within olive orchard (288).

The biggest number of *Coccinellidae* samples has been registered within olive groove (56), in comparison to orange groove (54).

ACKNOWLEDGMENTS

This research was accomplished within the doctoral program mobility from the University of Tras os Montes e Alto Douro of Vila Real, under the guidance of PhD Laura Monteiro Torres.

REFERENCES

- Andriev Sorina, Octavia, 2004** - Cercetări privind cunoașterea Coccinelidelor (Insecta-Coleoptera-Coccinellidae) din România din punct de vedere sistematic, biologic, ecologic și biogeografic Teza de doctorat, Universitatea „A.I.Cuza”, Iași, 294.
- Arion, G., 1912** - Raport asupra insectelor dăunătoare din familia Coccidelor. Buletin Agricol.;
- Baicu T., 1977** - Elaborarea măsurilor de combatere integrată. Probl. Prot. Plant.V.3; 203-221;
- Baicu, T., Săvescu, A., 1978** - Combaterea integrată în protecția plantelor - Editura Ceres, București;
- Baicu T., 1992** - Perspective în combaterea biologică a bolilor și dăunătorilor plantelor agricole, 72 p., Ed. Tehnica agricolă, București;
- Banița Emilia, Serafim Rodica, Searpe Doina, 1997** - Evoluția populațiilor de coccinele (Coleoptera Coccinellidae) în culturile de câmp din centrul Olteniei. Probl. de prot. Plant. XU (2);
- Baniță, E., Sterghiu, C., Luca E., Naidin, C., 1999** - Studiul paianjenilor prădători (Araneae) ai insectelor dăunătoare culturilor de cereale. Analele Institutului de Cercetări pentru Cereale și Plante Tehnice Fundulea, LXVI: 285-294.
- Bărbulescu, Al., 2001** - Rezultate obținute în anul 2000 în cadrul cercetărilor privind bolile și dăunătorii cerealelor privind bolile și dăunătorii cerealelor și unor plante tehnice și furajere. Probleme protecția plantelor, XXIX(2): 123-178.
- Boguleanu Gh., și colab., 1980** - Entomologie agricolă. Ed. Didactică și Pedagogică București;
- Borcea I., 1910** - Rolul insectelor prădătoare și parazite în agricultură. Rev. St. , V Adamachi. I, p.29-46, Iasi;
- Chapman and Hall, New York, USA Driesche RG, Hoddle MS, Center T., 2008** - Control of pests and weeds by natural enemies: an introduction to biological control. Wiley, New York;
- Ciepielowska D., 1991** - Ladybirds (Coleoptera-Coccinellidae) occurring on papilionaceous crops in the Olzyn District. Polskie Pismo Entomologiczne, G2 (1): 129-138;
- Ciochia V., 1986** - Combaterea biologică a dăunătorilor, verigă esențială a protecției ecosistemelor. Brașov
- Cozma V, Diaconu A., Grecu M., Tălmaciu M., Pareza Madalin, Vasiliu G., 2006** - Observații privind abundența și diversitatea coleopterelor din coronamentul unor livezi de mar cu management diferit de exploatare. Lucr. șt., Seria Horticultura, Iasi, I (49):1093-1096;
- Dejeu, L., Petrescu, C., Chira, A., 1997** - Horticultura și protecția mediului. Ed. Ceres, Bucuresti;
- Filip I. și colab., 2003**. Strategies of integrated control of the grape moth (*Lobesia botrana Den et Schiff.*) in Murfatlar vineyard. Analele Univ. din Craiova, Seria Biol., Horti., TPPA, vol VIII (XLIV) p.432.
- Ghizdavu I., Tomescu N., Oprean I., 1983**- Feromonii insectelor, "pesticide" din a III-a generație. Edit. Dacia, Cluj-Napoca.
- Ghizdavu L., Bunescu H., 1991**- Feromonii - important mijloc de combatere biotehnică a insectelor dăunătoare. Rev. Protecția Plantelor, nr.3-4. Tipografia Agronomică, Cluj-Napoca.
- Hodek L., 1973**- Biology of Coccinellidae. Academia Publishing House of the Czechoslovak Academy of Sciences, Prague, 260;
- Filip I. și colab., 2003** - Strategies of integrated control of the grape moth (*Lobesia botrana Den et Schiff.*) in Murfatlar vineyard. Analele Univ. din Craiova, Seria Biol., Horti., TPPA, vol VIII (XLIV) p.432.
- Foltz, J.L. 2002** - Coleoptera: Coccinellidae. Dept of Entomology and Nematology. University of Florida. http://entomology.ifas.ufl.edu/Coleoptera_Coccinellidae.html, modified 12/01/14.

- Ghizdavu I., Tomescu N., Oprean I., 1983 - Feromonii insectelor, "pesticide" din a III-a generație.** Edit. Dacia, Cluj-Napoca.
- Hodek L., 1973 - Biology of Coccinellidae.** Academia Publishing House of the Czechoslovak Academy of Sciences, Prague, 260;
- Iperti, G., 1974 - Les principaux auxiliaires entomophages Coleopteres. Les organismes auxiliaires en vergers de pommieres.** Brochure Bulletin International Biological Prevention Organization, West Palearctic Regional Section: 111-121 ;
- Isac Gr., 1989- Biotehnologia combaterii molilor strugurilor prin folosirea entomofagilor.** Lucr. St. ICVV Ștefănești Argeș, p 279-292. București 1989.
- Kalushkov P. K., Denkov K.D., Dimova V., I., 1990- Ecofaunistic studies on the coccinellids (Coleoptera-Coccinellidae) in lucerne plats near Pleven, Ekologiya., Bulgaria.** Zool. Inst. Bulgarien Acad. Sci.23: 37-66;
- Klausnitzer,B., Klausnitzer H., 1983 - Marienkäfer (Coccinellidae),** Amandus Ziemsen Verlag, Wittenberg, 103 pp;
- Klausnitzer, B 2004- *Harmonia axyridis* (Pallas, 1773) in Bassel-Stadt (Coleoptera Coccinellidae).** Mitteilungen Entomol. Gesellschaft Basel54: 115-122;
- Kovalenkov V.G., Tyurina N.M., 1992- Coccinellids against alfaifa pest.** Zashchita Rasteni (Moskva), 10; 13-14;
- Lăcătușu Matilda, Tudor Constanța, Teodorescu Irina Nadej. M, 1981- Structura faunistică din culturi de lucernă.** Stud. Cerc. Biol. Anim, 33, 2: 179-182;
- Malschi Dana, Mustea D., 1992- Dinamica entomofaunei dăunătoare specifice agrobiocenozelor de grâu din centrul Transilvaniei în perioada 1981-1990.** Probl. Prot. Plantelor, vol. XX nr.3-4, 237-348;
- Sonia A. P. Santos., 2012 - Species abundance patterns of coccinellid communities associated with olive, chestnut and almond crops in north-eastern Portugal,** Agricultural and Forestry Entomology, 14(4) DOI: 10.1111/j.1461-9563.2012.00578.x, 4-6.