

## THE POTENTIAL OF SOME INDIGENOUS MICROORGANISMS AND PLANTS FOR THE REMOVAL OF HEAVY METALS FROM SOIL

### POTENȚIALUL UNOR MICROORGANISME ȘI PLANTE INDIGENE DE ELIMINARE A METALELOR GRELE DIN SOL

ROȘCA Mihaela<sup>1</sup>, HLIHOR Raluca-Maria<sup>1,2</sup>, COMĂNIȚĂ Elena-Diana<sup>1</sup>, COZMA Petronela<sup>1</sup>, SIMION Isabela-Maria<sup>1,2</sup>, GAVRILESCU Maria<sup>1,3</sup>,  
e-mail: raluca.hlihor@uaiasi.ro; mgav@ch.tuiasi.ro

**Abstract.** Heavy metals found in soils from different industrial sources or mining activities are persistent inorganic pollutants able to bioaccumulate along the food chain and cause negative effects in the environment and for human health. Different physical, chemical and biological processes are applied for their removal from soil environments. Biological processes become more and more preferred, since bioremediation strategies have often proved to be more advantageous than the conventional remediation tools, mainly because these processes can be implemented directly onto the contaminated sites (in situ). In this context, the present paper examines the ability of microorganisms and plants to remove heavy metals from soil, in terms of tolerance and bioaccumulation. A particular interest is given to the bioaccumulation processes of metals by proteobacteria, bacilli and actinobacteria, alone or in synergism with indigenous plants. Also, some advances in the biosorption of highly toxic heavy metal ions as Cr(VI) and Cd(II) are just discussed, together with various strategies and practices to explore the synergism between microorganisms and plants as valuable biological resource for increasing tolerance against heavy metals and strengthening the bioremediation processes. **Key words:** bioaccumulation, microorganisms, plants, soil pollution, synergism, tolerance

**Rezumat.** Metalele grele provenite din diferite surse industriale sau din activitățile miniere sunt poluanți anorganici persistenti ai solurilor, capabili să se bioacumuleze de-a lungul lanțului trofic și să genereze efecte negative pentru mediu și sănătatea umană. Pentru îndepărtarea metalelor grele din sol se pot aplica procese fizice, chimice și biologice. Procesele biologice sunt preferate din ce în ce mai mult, deoarece strategiile de bioremediere s-au dovedit adesea mai avantajoase decât instrumentele de remediere convenționale și pot fi implementate direct pe siturile contaminate (in situ). În acest context, lucrarea analizează capacitatea unor microorganisme și plante indigene de a elimina metalele grele din sol, în relație cu aspecte ce privesc toleranța și bioacumularea metalelor grele. O atenție aparte se acordă proceselor de bioacumulare a metalelor în proteobacterii, bacili și actinobacterii, singure sau

---

<sup>1</sup>“Gheorghe Asachi” Technical University of Iași, Romania

<sup>2</sup>University of Agricultural Sciences and Veterinary Medicine Iași, Romania

<sup>3</sup>Academy of Romanian Scientists, Bucharest, Romania

*în sinergism cu unele plante indigene. De asemenea sunt evidențiate progresele în biosorbția ionilor unor metale grele foarte toxice, precum Cr(VI) și Cd(II) și sunt discutate pe scurt diferite strategii și practici destinate explorării sinergismului dintre microorganismele și plante ca o resursă biologică valoroasă pentru creșterea toleranței la metale grele și progresului în cunoașterea proceselor de bioremediere.*

**Cuvinte cheie:** bioacumulare, microorganismele, plante, poluarea solului, sinergism, toleranță

## INTRODUCTION

Nowadays numerous sites worldwide are contaminated with various organic and inorganic pollutants having different toxic and persistence characteristics. The presence of inorganic toxic pollutants in soils is especially given by heavy metals ions (Gavrilăscu, 2014; Pavel *et al.*, 2012; Sobariu *et al.*, 2017; Tóth *et al.*, 2016a). This type of pollution can cause negative impacts in the environment and for ecological systems (e.g. inhibition of cytoplasmic enzymes, damage to cell structures, inhibition of protein synthesis, negative effects in animals by disturbing the central and peripheral nervous and circulatory systems, decrease of plants growth, performance and crop yield etc.) (Cozma and Gavrilăscu, 2014; Hlihor *et al.*, 2017; Siminciuc *et al.*, 2015; Sobariu *et al.*, 2017). As a consequence, human health can suffer numerous injuries (e.g. on the functions of brain, lungs, kidney, liver, blood composition; physical, muscular, and neurological degenerative diseases; immune deficiency disorders, heart complications, digestion problems, cancer etc.) (Hlihor *et al.*, 2017; lordache *et al.*, 2016; Rosca *et al.*, 2015; Sobariu *et al.*, 2017).

Based on the information provided by LUCAS database, Tóth *et al.* (2016a) found that about 1.2 million km<sup>2</sup> (28.3%) of the total surface area of the European Union are affected by pollution with one or more heavy metals ions (e.g. As, Cd, Cr, Cu, Pb, Zn, Sb, Co, Ni), whose concentration exceeds the maximum admissible limits. The largest areas affected by this type of pollution are situated in the West and Central of Europe, center of Italy, Greece and South-East of Ireland. The highest concentrations of Pb in soils were found in the center of Italy, France, Germany, and United Kingdom. Ni is mostly abundant in Mediterranean region of Europe, especially in Greece, while pollution with chromium affects largest areas of Piemonte, Lorraine-Alsace, Western-Macedonia and center of Greece. High concentrations of cadmium can be found in Ireland and Greece, whereas the concentration of Hg ions exceeds the legal limits especially in the center of Italy, North-West of England and East of Slovakia (Tóth *et al.*, 2016a, 2016b). In Romania, approximately 6,639 ha are affected by heavy metals pollution, noting that for 5,773 ha the concentration of some heavy metals found in soil exceeds the maximum admissible limits. The most affected areas are in the West of Romania (23.2%), North - East (20.5%), North - West (19.7%), Center (12.3%) and South - West Oltenia (2%) (Dumitru *et al.*, 2011).

Various physical, chemical and biological processes can be applied for the decontamination of soils polluted with heavy metals, but a special importance is given today to bioremediation techniques (fig. 1). Biological processes become more and more preferred, since the bioremediation strategies have often proved to be more advantageous than the conventional remediation tools, mainly because they are economic and ecological friendly and can be implemented directly onto the contaminated sites (*in situ*). Heavy metals can be eliminated by their bioaccumulation in specific microorganisms and plants. In this context, the present paper examines the abilities of some microorganisms and plants to eliminate heavy metals from polluted soils, considering also their tolerance and bioaccumulation capacity for heavy metals.




Fig. 1 Sources of heavy metals in soils and the processes used for remediation

## BIOREMEDIATION OF SOILS

Recent studies have shown that biological methods based on phytoremediation and bioaccumulation are two *in situ* eco-friendly and economically feasible methods applied for an efficient soil remediation. These processes are based on the potential of plants (native, woody, herbaceous, ornamental, perennial, flowering, wild etc.) or microorganisms (bacteria, fungi, yeasts) to remove heavy metal ions from soil by extracting, transferring and accumulating them in biomass (Colin *et al.*, 2012; Kumari *et al.*, 2016; Rosca *et al.*, 2015; Sobariu *et al.*, 2017). The results of some studies on the potential of plants (tab. 1) and microorganisms (tab. 2) to eliminate heavy metals from soils by their bioaccumulation have demonstrated that various plants and microorganisms possess a high capacity to tolerate and absorb heavy metal ions, playing also an important role in reducing the contamination risk of food with toxic metals (Hlihor *et al.*, 2017; Sobariu *et al.*, 2017).

## Removal of heavy metals ions by plants (phytoremediation)

Species of plants	Removal capacity of heavy metals ions (mg/kg d.w*.)								References
	Fe <sup>2+</sup>	Pb <sup>2+</sup>	Ni <sup>2+</sup>	Cu <sup>2+</sup>	Zn <sup>2+</sup>	Cd <sup>2+</sup>	Cr <sub>tot</sub>	Co <sup>2+</sup>	
<i>Brassica napus</i>	-	472	45.7	414	5983	23.5	82.5	-	Marchiol <i>et al.</i> (2004)
<i>Raphanus sativus</i>	-	407	51.6	563	4029	59.5	62	-	
<i>Cardaria draba</i>	1452	827	9.5	34.4	1850	2.2	-	-	Chehregani <i>et al.</i> (2009)
<i>Amaranthus retroflexus</i>	6230	371.5	7	57.5	233	5.5	-	-	
<i>Boromus sp</i>	520	210	6.5	26	85	10	-	-	
<i>Noea mucronata</i>	1230	1485	18.5	84	1984	14.6	-	-	
<i>Marrubium vulgare</i>	540	78	4	34	58	9	-	-	Porębska and Ostrowska (1999)
<i>Lactuca serriola</i>	-	3	8	18	1030	21	4	-	
<i>Artemisia vulgaris</i>	-	17	6.8	81	398	36	4	-	Ipătioaie <i>et al.</i> (2014)
<i>Rheum rhabarbarum L.</i>	-	0.015	-	0.03	0.17	-	-	-	Shallari <i>et al.</i> (1998)
<i>Herniaria hirsuta</i>	-	34	808	22	29	8	275	63	
<i>Inula germanica</i>	-	24	211	20	24	5	89	31	
<i>Dittrichia graveolens</i>	-	28	94	1110	849	9	69	34	
<i>Lotus ormithopodioides</i>	-	43	232	14	75	5	63	21	
<i>Alyssum murale</i>	-	23	8463	23	108	3	12	86	
<i>Convolvulus arvensis L.</i>	-	-	-	560	-	1500	800	-	Gardea-Torresdey <i>et al.</i> (2004)
<i>Euphorbia cheiradenia</i>	1040	1138	14.2	65	1873	2.35	-	-	Chehregani and Malayeri (2007)
<i>Biebers teiniamultifida</i>	480	23	4	20	-	7	-	-	
<i>Reseda lutea</i>	5490	371	7	57.5	233	5.5	-	-	
<i>Euphorbia macroclada</i>	2261	81.67	13	26	327	3	-	-	Matache <i>et al.</i> (2013)
<i>Ceratophyllum demersum</i>	-	20	-	22.7	104	3.52	-	-	
<i>Potamogeton pectinatus</i>	-	6.63	-	6.24	16.4	0.64	-	-	
<i>Potamogeton lucens</i>	-	1.51	-	9.8	15.6	0.97	-	-	

\*d.w. – dry weight

Phytoremediation and bioaccumulation processes applied in contaminated areas can be influenced by several operational parameters (soil pH, temperature variations, soil morphology, initial concentration of heavy metals etc.) (Pavel *et al.*, 2013). Taking into account these aspects, the selection of the most suitable plants and microorganisms for bioremediation depends on their ability to adapt to the climatic and soil conditions in the polluted area.

## Removal of heavy metals ions by microorganisms through bioaccumulation process

Microorganisms species	Initial conditions	Ions	Removal efficiency (%) or uptake (mg/g)	References
<i>Arthrobacter phenanthrenivorans</i>	$C_{iPb(II)}=180$ mg/L, $C_{iCd(II)}=178$ mg/L, $C_{iNi(II)}=85$ mg/L, 72 hours contact time	Pb <sup>2+</sup>	79.9 %	Banerjee et al. (2016)
		Ni <sup>2+</sup>	47.62%	
		Cd <sup>2+</sup>	34.05%	
<i>Pseudomonas putida</i>	$C_{iPb(II)}=1.53$ mg/L, $C_{iCo(II)}=8.82$ mg/L, $C_{iNi(II)}=11.48$ mg/L, $C_{iCu(II)}=5.9$ mg/L, $C_{iCd(II)}=50$ mg/L, 5 days contact time	Pb <sup>2+</sup>	96%	Nanganuru and Korrapati (2012)
		Ni <sup>2+</sup>	57%	
		Cu <sup>2+</sup>	49 %	
		Co <sup>2+</sup>	71%	
		Cd <sup>2+</sup>	93%	
<i>Enterobacter</i> sp. J1	$C_{iheavy\ metal}=100$ mg/L, 24 hours contact time	Pb <sup>2+</sup>	50.9 mg/g	Lu et al. (2006)
		Cu <sup>2+</sup>	32.5 mg/g	
		Cd <sup>2+</sup>	46.6 mg/g	
<i>Bacillus cereus</i>	$C_{iheavy\ metal}=0.2-0.6$ mmol/mL, 24 hours contact time	Pb <sup>2+</sup>	36 mg/g	Banerjee et al. (2015)
		Ni <sup>2+</sup>	28 mg/g	
		Cu <sup>2+</sup>	33 mg/g	
		Mn <sup>2+</sup>	38 mg/g	
		Hg <sup>2+</sup>	35 mg/g	
		Co <sup>2+</sup>	31 mg/g	
<i>Saccharomyces cerevisiae</i>	$C_{iHg(II)}, Co(II)}=500$ mg/L, 21 days contact time $C_{iPb(II)}=100$ mg/L $C_{iCd(II)}=300$ mg/L 30 days contact time	Hg <sup>2+</sup>	90.48 %	Imam et al. (2016)
		Cd <sup>2+</sup>	92.68 %	Damodaran et al. (2011)
		Pb <sup>2+</sup>	67 %	
		Cd <sup>2+</sup>	73 %	
<i>Bacillus subtilis</i> B38	$C_{iheavy\ metal}=0.1-1.25$ mmol/L 24 hours contact time	Pb <sup>2+</sup>	2.48 mmol/g	Wang et al. (2014)
		Hg <sup>2+</sup>	4.09 mmol/g	
		Cd <sup>2+</sup>	3.04 mmol/g	
		Cr <sub>total</sub>	1.83 mmol/g	
Mix of proteobacteria: <i>O. intermedium</i> , <i>A. ebreus</i> , <i>A. caviae</i> , <i>B. diminuta</i> , <i>B. vietnamiensis</i> , <i>P. mendocina</i> , <i>P. alcaligenes</i> , <i>S. marcescens</i> , <i>S. acidaminiphilia</i> , <i>D. tsuruhatensis</i>	$C_{iAs(II)}=141$ mg/kg soil, 20 min contact time $C_{iZn(II)}=49$ mg/kg soil $C_{iNi(II)}=21$ mg/kg soil 10 min contact time	As <sup>2+</sup>	71%	Fauziah et al. (2017)
		Ni <sup>2+</sup>	50.8%	
		Zn <sup>2+</sup>	47.6%	
<i>Thiobacillus ferrooxida</i>	$C_i=2199$ mg/kg soil $C_i=27660$ mg/kg soil $C_i=132$ mg/kg soil $C_i=99500$ mg/kg soil $C_i=7520$ mg/kg soil $C_i=11.68$ mg/kg soil $C_i=7360$ mg/kg soil 72 hours contact time	Zn <sup>2+</sup>	90 %	Sur et al. (2012)
		Pb <sup>2+</sup>	0.53 %	
		Cr <sub>total</sub>	25 %	
		Fe <sup>2+</sup>	100 %	
		Cu <sup>2+</sup>	100 %	
		Cd <sup>2+</sup>	60 %	
		Mn <sup>2+</sup>	65 %	

In this case, the best choices for soil bioremediation is through plants that normally grow in the contaminated area, as well as indigenous microorganisms. Plants and microorganisms age and the synergistic or antagonistic effects of elements/compounds on going in soil against plants and microorganisms are also important factors that can influence process efficiency (Elekes, 2014). Barbeș and

Bărbulescu (2017) analyzed the accumulation of Cu, Zn, Ni, Pb, Cr, Cd and Co in the leaves and bark of *Populus nigra* L., a species of plant grown in Navodari area, in the region of Romanian Black Sea Littoral. The results have shown that the plant possesses a high potential in the removal of heavy metals ions from polluted area (the effectiveness of the process was higher than 70%). Elekes (2014) studied the potential of some plants, which normally grow on the industrial platform of Targoviste city (*Lolium perenne*, *Festuca pratensis*, *Stipa capillata*, *Agrostis alba*, *Cynodon dactylon*, *Agrostis tenuis* and *Luzula campestris*). The samples taken were analyzed and the results have shown that the plants studied are able to remove the heavy metals (Cu, Zn, Sn, Pb, Co, Ni, Mn, Cr) (e.g. *Lolium perenne* has removed 921.67 mg/kg Zn, 201.23 mg/kg Pb, 114.19 mg/kg Cr, 61.95 mg/kg Cu).

## CONCLUSIONS

When comparing the data available from literature, we can notice that biological *in situ* processes can be applied with high potential for the removal of heavy metals from polluted soils, considering different plants and microorganisms. Some species of plants such as: *Amaranthus retroflexus*, *Noea mucronata*, *Alyssum murale*, *Convolvulus arvensis* L., *Brassica napus*, *Raphanus sativus* and *Cardaria draba* can be considered as hyperaccumulators for some heavy metals ions. Also, it has been observed that a wide variety of indigenous microorganisms from proteobacteria, actinobacteria and bacilli class can be used with high effectiveness for the bioremediation of soils polluted with heavy metals (e.g.: *S. cerevisiae*, *Streptomyces*, *A. phenanthrenivorans*, *Pseudomonas putida*, *Enterobacter* sp. etc.). In Romania, the use of these processes for the remediation of soils polluted with various heavy metal ions can be successfully applied, as confirmed by the studies carried out in Navadori and Targoviste regions, as well as in other areas of the country. In order to increase the efficiency of the process, it is necessary to select plants and microorganisms adapted or easily adaptable to climatic conditions in the area, structure and parameters of the polluted soil.

**Acknowledgments:** This work was supported by a grant of the Romanian National Authority for Scientific Research, CNCS – UEFISCDI, project number PN-III-P4-ID-PCE-2016-0683, Contract no. 65/2017.

## REFERENCES

1. Banerjee S., Gothalwal R., Sahu P.K., Sao S., 2015 - *Microbial Observation in Bioaccumulation of Heavy Metals from the Ash Dyke of Thermal Power Plants of Chhattisgarh, India*. Advances in Bioscience and Biotechnology, 2015, 6, 131-138.
2. Banerjee G., Ray A.K., Pandey S., Kumar R., 2016 - *An Alternative Approach of Toxic Heavy Metal Removal by Arthrobacter phenanthrenivorans: Assessment of Surfactant Production and Oxidative Stress*. Current Science, 110, p. 2124-2128.

3. **Barbeș L., Bărbulescu A., 2017** -*Monitoring and Statistical Assessment of Heavy Metals in Soil and Leaves of Populus nigra L.* Environmental Engineering and Management Journal, 16, p. 187-196.
4. **Chehregani A., Malayeri B.E., 2007** - *Removal of Heavy Metals by Native Accumulator Plants.* International Journal of Agriculture & Biology, 9, p. 462-465.
5. **Chehregani A., Noori M., Yazdi H.L., 2009** - *Phytoremediation of Heavy-Metal-Polluted Soils: Screening for New Accumulator Plants in Angouran Mine (Iran) and Evaluation of Removalability.* Ecotoxicology and Environmental Safety, 72, p. 1349-1353.
6. **Colin V.L., Villegas L.B., Abate C.M., 2012** - *Review: Indigenous Microorganisms as Potential Bioremediators for Environments Contaminated with Heavy Metals.* International Biodeterioration & Biodegradation, 69, 28-37.
7. **Cozma P., Gavrilesu M., 2014** - *Key Features of Contaminated Soils, In: Contaminated Soils: a Guide to Sampling and Analysis.* Gavrilesu M., Bulgariu L. (Eds), Future Science, London, p. 7-17.
5. **Damodaran D., Suresh G., Mohan R.B., 2011** -*Bioremediation of Soil by Removing Heavy Metals Using Saccharomyces cerevisiae.* 2<sup>nd</sup> International Conference on Environmental Science and Technology IPCBEE, 6, p. V2-22-V2-27.
6. **Dumitru M., Manea A., Ciobanu C., Dumitru S., Vrinceanu N., Rișnoveanu I., Calciu I., Tănase V., Preda M., Mocanu V., Eftene M., 2011** - *Soil Quality Monitoring In Romania.* Editura SITECH, Craiova, Romania.
7. **Elekes C.C., 2014** - *Chapter 10: Eco-Technological Solutions for the Remediation of Polluted Soil and Heavy Metal Recovery, In: Environmental Risk Assessment of Soil Contamination.* InTechEditure, p. 309-335.
8. **Fauziah S.H., Jayanthi B., Emenike C.U., Agamuthu, 2017** -*Remediation of Heavy Metal Contaminated Soil Using Potential Microbes Isolated from a Closed Disposal Site,* International Journal of Bioscience, Biochemistry and Bioinformatics, 7, p. 230-237.
9. **Gardea-Torresdey J.L., Peralta-Videa J.R., Montes M., de la Rosa G., Corral-Diaz B., 2004** - *Bioaccumulation of Cadmium, Chromium and Copper by Convolvulus arvensis L.: Impact on Plant Growth and Uptake of Nutritional Elements.* Bioresource Technology, 92, p. 229-235.
10. **Gavrilesu M., 2014** - *Colloid-Mediated Transport and the Fate of Contaminants in Soils, In: The Role of Colloidal Systems in Environmental Protection.* Elsevier, Amstredam, p.397-451.
11. **Hlihor R.-M., Apostol L.C., Gavrilesu M., 2017** - *Environmental Bioremediation by Biosorption and Bioaccumulation: Principles and Applications, In: Enhancing Cleanup of Environmental Pollutants, Volume 1: Biological Approaches.* Anjum N., Gill S., Tuteja N., (Ed.), Springer Science+Business Media, New York, p. 289-315.
12. **Imam A.S.S., Rajpoot I.K., Gajjar B., Sachdeva A., 2016** - *Comparative Study of Heavy Metal Bioremediation in Soil by Bacillus Subtilis and Saccharomyces erevisiae.* Indian Journal of Science and Technology, 9, p. 1-7.
13. **Iordache M., Branzoi I.V., Popescu L.R., Iordache I., 2016** - *Evaluation of Heavy Metal Pollution into a Complex Industrial Area From Romania.* Environmental Engineering and Management Journal, 15, 389-394.
14. **Ipățioaie D.C., Munteanu N., Stoleru V., Sellitto V.M., Cojocaru A, 2014** – *The accumulation of heavy metals in rhubarb (Rheum rhabarbarum L.).* Lucrări științifice, seria Horticultură, 57, p. 277-282.
15. **Kumari A., Lal B., Rai U.N., 2016** - *Assessment of Native Plant Species for Phytoremediation of Heavy Metals Growing in the Vicinity of NTPC Sites, Kahalgaon, India.* International Journal of Phytoremediation, 18, 592-597.

16. Lu W.B., Shi J.J., Wang C.H., Chang J.S., 2006 - *Biosorption of Lead, Copper and Cadmium by an Indigenous Isolate Enterobacter sp. J1 Possessing High Heavy-Metal Resistance*. Journal of Hazardous Materials, 134, 80-86.
17. Marchiol L., Assolari S., Sacco P., Zerbi G., 2004 - *Phytoextraction of Heavy Metals by Canola (Brassica napus) and Radish (Raphanussativus) Grown on Multicontaminated Soil*. Environmental Pollution, 132, p. 21-27.
18. Matache M.L., Marin C., Rozyłowicz L., Tudorache A., 2013 - *Plants accumulating heavy metals in the Danube River wetlands*. Journal of Environmental Health Science and Engineering, 11, p. 1-7.
19. Nanganuru H.Y., Korrapati N., 2012 - *Studies on Biosorption of Cadmium by Pseudomonasputida*. International Journal of Engineering Research and Applications, 2, p. 2217-2219.
20. Pavel V.L., Stătescu F., Gavrilescu M., 2012 - *Aspects Concerning the Behaviour of Heavy Metals in Soil*, In: *Current topics, Concepts and Research Priorities in Environmental Chemistry*. Zaharia C. (Ed.), Editura Universității „Alexandru Ioan Cuza” din Iași, pp. 341-358.
21. Pavel V.L., Sobariu D.-L., TudoracheFertu I.D., Stătescu F., Gavrilescu M., 2013 - *Symbiosis in the Environment: Biomangementof Soils Contaminated with Heavy Metals*. European Journal of Science and Theology, 9, 211-224.
22. Rosca M., Hlihor R.-M., Cozma P., Comăniță E.-D., Simion I.M., Gavrilescu M., 2015 - *Potential of biosorption and bioaccumulation processes for heavy metals removal in bioreactors*. E-Health and Bioengineering Conference (EHB), DOI: 10.1109/EHB.2015.7391487.
23. Shallari S., Schwartz C., Haskoba A., Morela J.L., 1998 - *Heavy Metals in Soils and Plants of Serpentine and Industrial Sites of Albania*. The Science of the Total Environment, 209, p. 133-142.
24. Siminciuc C.G., Munteanu N., Teliban G., Stoleru V., 2015 - *Assement of heavy metals content on tomatoes under organic system*. Lucrări științifice, seria Horticultură, USAMV Iași, 58, 279-284.
25. Sobariu D.L., TudoracheFertu D.I., Diaconu M., Pavel L.V., Hlihor R.-M., Drăgoi E.N., Curteanu S., Lenzde M., Corvini P.F.X., Gavrilescu M., 2017 - *Rhizobacteria and Plant Symbiosis in Heavy Metal Uptake and Its Implications for Soil Bioremediation*. New Biotechnology, 39, p. 125-134.
26. Sur I.M.B., Micle V., Avram S., Șenilă M., Oros V., 2012 - *Bioremediation of Some Heavy Metals from Polluted Soils*. Environmental Engineering and Management Journal, 11, p. 1389-1393.
27. Tóth G., Hermann T., Szatmári G., Pásztor L., 2016a - *Maps of Heavy Metals in the Soils of the European Union and Proposed Priority Areas for Detailed Assessment*. Science of the Total Environment, 565, p. 1054-1062.
28. Tóth G., Hermann T., Da Silva M.R., Montanarella L., 2016b - *Heavy Metals in Agricultural Soils of the European Union with Implications for Food Safety*. Environment International, 88, p. 299-309.
29. Wang T., Sun H., Mao H., Zhang Y., Wang C., Zhang Z., Wang B., Sun L., 2014 - *The Immobilization of Heavy metals in Soil by Bioaugmentation of aUV-mutant Bacillus subtilis 38 Assisted by NovoGroBiostimulation and Changes of Soil Microbial Community*. Journal of Hazardous Materials, 278, p. 483-490.